



# CJC-1295 (NO DAC) / IPAMORELIN (Blend)

PEPTIDE RESEARCH COMPOUND

## **Dilution Guidelines**

- Use **1.5–2 mL BAC water** per 5 mg / 5 mg blend
- Add BAC water **slowly along the inner wall** of the vial—**do not inject directly onto the powder**
- Allow both vial and BAC water to reach room temperature before reconstitution
- A slightly higher volume of BAC water may help if the solution appears too thick

## **Appearance After Mixing**

- Solution typically appears **clear to mildly cloudy**
- **Swirl or roll gently**—do not shake, as both peptides are moderately sensitive to agitation
- Allow a few minutes for **complete dissolution** before use

## **Storage**

- Store at **2–8°C (refrigerated)** after mixing
- Do not freeze and keep protected from **light and air**
- Use within **10–14 days**

## **Areas of Research Interest:**

- Studied for their **complementary activity** in stimulating natural growth hormone release
- Frequently explored in models involving **muscle preservation, tissue regeneration, and cellular repair**
- Investigated for **short-acting GH pulse modulation** with **minimal cortisol or prolactin stimulation**
- Considered in age-related studies focused on **hormonal support, recovery, and metabolic stability**
- Used in research comparing **GHRH analogs** and **ghrelin mimetics** in combination protocols

## **DISCLAIMER**

This compound is provided for laboratory research purposes only. It is not intended for human or veterinary use, and no therapeutic or diagnostic claims are made.