



# CJC-1295 (WITH DAC) / / IPAMORELIN (Blend)

PEPTIDE RESEARCH COMPOUND

## **Dilution Guidelines**

- Use **1.5–2 mL BAC water** per 5 mg / 5 mg blend
- Add BAC water **slowly along the inner wall** of the vial—**do not inject directly onto the lyophilized blend**
- Allow both the vial and BAC water to reach **room temperature** before mixing
- Slightly more fluid may be used if the solution appears **dense or slow to draw**

## **Appearance After Mixing**

- Typically forms a **clear to faintly cloudy** solution
- **Swirl or roll gently** to dissolve—**do not shake**, as the DAC-modified structure is more sensitive to agitation
- Allow several minutes for **full dissolution** before use

## **Storage**

- Store at **2–8°C (refrigerated)** after reconstitution
- Do not freeze
- Use within 15–20 days
- Keep tightly sealed and protect from light and air between uses

## **Areas of Research Interest:**

- Studied for their **combined action** on **sustained** (DAC) and **pulsatile** (Ipamorelin) growth hormone release
- Frequently explored in **long-term research models** related to muscle regeneration, metabolic function, and tissue repair
- Investigated for **extended half-life, enhanced IGF-1 activity**, and **receptor pathway synergy**
- Considered in age-related studies focused on **hormonal optimization, body composition**, and **fat metabolism**
- Used to assess the interaction of **GHRH analogs** and **ghrelin receptor agonists** in experimental endocrine protocols

## **DISCLAIMER**

This compound is provided for laboratory research purposes only. It is not intended for human or veterinary use, and no therapeutic or diagnostic claims are made.