



CJC-1295 (WITH DAC)

PEPTIDE RESEARCH COMPOUND

Dilution Guidelines

- Minimally 1–1.5 mL BAC water per 5 mg vial
- Add BAC water slowly along the inner wall—avoid injecting directly onto the peptide
- Allow vial and BAC water to reach room temperature before reconstitution
- Swirl gently—do not shake vigorously, as this may affect peptide structure

Appearance After Mixing

- Typically results in a clear to slightly cloudy solution
- Mild foaming may occur—allow to settle and swirl gently again if needed
- Full dissolution generally occurs within 1–2 minutes

Storage

- Store at **2–8°C (refrigerated)** after reconstitution
- Do not freeze and protect from light and air exposure
- Use within 10–21 days for optimal stability

Areas of Research Interest:

- Researched for its ability to extend **growth hormone release** over a prolonged period
- Frequently explored in long-term models related to **muscle growth, body composition, and endocrine balance**
- Studied for consistent stimulation of **IGF-1 production** across multiple days
- Compared to non-DAC versions for its impact on **hormonal stability and dosing intervals**
- Investigated in studies involving **metabolic function, tissue repair, and age-related decline**

DISCLAIMER

This compound is provided for laboratory research purposes only. It is not intended for human or veterinary use, and no therapeutic or diagnostic claims are made.

Note:

CJC-1295 is available in two research formats: with or without **DAC (Drug Affinity Complex)**. The DAC version is specifically studied for its **extended half-life and prolonged physiological interaction** compared to the shorter-acting non-DAC variant.