



EPITHALON

PEPTIDE RESEARCH COMPOUND

Dilution Guidelines

- Minimally 1.5–2 mL BAC water per 5 mg vial
- Tilt the vial and allow water to run slowly along the inner wall—avoid injecting directly onto the powder
- Add BAC water gradually to minimize foaming and ensure even reconstitution
- Swirl or roll gently—do not shake, as Epithalon is moderately delicate

Appearance After Mixing

- Typically forms a clear or slightly hazy solution
- Light bubbling or foaming may occur; let it settle naturally before use
- Full dissolution should occur within 1–2 minutes at room temperature

Storage

- Store at **2–8°C (refrigerated)** after reconstitution
- Do not freeze and protect from light and air exposure
- Keep upright and sealed when not in use
- Use within 10–21 days for optimal stability

Areas of Research Interest:

- Frequently studied for its potential influence on **cellular aging** and **telomere support**
- Investigated in research related to **pineal gland function** and **melatonin regulation**
- Explored in models addressing **oxidative stress**, **DNA repair**, and **immune function**
- Considered in aging-related studies focused on **biological markers** and **longevity**
- Used in studies involving **circadian rhythm regulation** and **hormonal balance**

DISCLAIMER

This compound is provided for laboratory research purposes only. It is not intended for human or veterinary use, and no therapeutic or diagnostic claims are made.